Quinine blocks 5-HT and 5-HT $_3$ receptor mediated peristalsis in both guinea pig and mouse ileum tissue

Jacqueline Walsh¹, Iain Barrett¹, Andrew Thompson², Sarah Lummis², Stephen Kelley¹. ¹University of Kent, Chatham Maritime, UK, ²University of Cambridge, Cambridge, UK.

Introduction. Quinine is commonly used to treat malaria; however one of the principal side effects is gastrointestinal disturbances (White, 1992). $5-HT_3$ receptors modulate gut peristalsis (Chetty *et al.,* 2006), and, as quinine has been shown to act as a $5-HT_3$ receptor antagonist (Thompson and Lummis, 2008) it is possible that these side effects result from actions at gut $5-HT_3$ receptors. To address this question, we examined the ability of quinine to antagonise 5-HT and $5-HT_3$ mediated peristalsis in guinea pig and mouse ileum.

<u>Methods</u>. Ileum was excised from male guinea pigs (200-300g) and C57BL/6 mice (25-35g) following cervical dislocation. Ileum segments (3-5 cm) were mounted in 50 ml organ baths containing Tryode's solution at 35-37 °C. Concentration-response curves were constructed for 5-HT and the selective 5-HT₃ agonist 2-Me-5-HT (non-cumulative doses). Quinine was pre-applied for 10 min and inhibition measured using agonist concentrations that elicited a submaximal response.

<u>Results</u>. Concentration-dependent contractions produced by 5-HT ($pEC_{50} = 5.45 \pm 0.17$, n = 8) and the selective 5-HT₃ agonist 2-Me-5-HT (5.01 ± 0.17, n = 11) were not significantly different (Student's *t*-test, t = 0.619, df = 17, p = 0.544) in guinea pig ileum. Increasing concentrations of quinine were able to antagonise the activities of both 5-HT ($p|C_{50} = 5.03 \pm 0.2$, n = 6) and 2-Me-5HT ($p|C_{50} = 4.59 \pm 0.26$, n = 4). At mouse ileum, 5-HT ($pEC_{50} = 7.57 \pm 0.33$, n = 9) was more potent (Student's *t*-test, t = 3.6, *df* = 12, p = 0.004) than 2-Me-5-HT ($pEC_{50} = 5.45 \pm 0.58$, n = 5). Quinine antagonised both the 5-HT ($p|C_{50} = 4.87 \pm 0.31$, n = 7) and 2-Me-5-HT-induced ($p|C_{50} = 6.18 \pm 1.14$, n = 4) contractions.

<u>Conclusions</u>. These results support previous electrophysiological studies that identified quinine as an antagonist at recombinant 5-HT₃ receptors with IC_{50} values comparable with those reported here (p/ C_{50} = 4.87, Thompson *et al.*, 2007). Further, we found that quinine completely blocked 5-HT induced contractions in mouse and guinea pig, raising the possibility that quinine targets other 5-HT receptors in the gut (e.g., 5-HT₄ receptors) and may influence intestinal function.

Chetty *et al.*, (2006). *Br.J.Pharmacol.*, 148; 112-141. Thompson *et al.*, (2007). *Br.J.Pharmacol* 151; 666-667. Thompson and Lummis, (2008). *Br.J.Pharmacol* 153; 1686-96. White, (1992). *BJCP*, 34; 1-10.